



This is a digital copy of a book that was preserved for generations on library shelves before it was carefully scanned by Google as part of a project to make the world's books discoverable online.

It has survived long enough for the copyright to expire and the book to enter the public domain. A public domain book is one that was never subject to copyright or whose legal copyright term has expired. Whether a book is in the public domain may vary country to country. Public domain books are our gateways to the past, representing a wealth of history, culture and knowledge that's often difficult to discover.

Marks, notations and other marginalia present in the original volume will appear in this file - a reminder of this book's long journey from the publisher to a library and finally to you.

Usage guidelines

Google is proud to partner with libraries to digitize public domain materials and make them widely accessible. Public domain books belong to the public and we are merely their custodians. Nevertheless, this work is expensive, so in order to keep providing this resource, we have taken steps to prevent abuse by commercial parties, including placing technical restrictions on automated querying.


We also ask that you:

- + *Make non-commercial use of the files* We designed Google Book Search for use by individuals, and we request that you use these files for personal, non-commercial purposes.
- + *Refrain from automated querying* Do not send automated queries of any sort to Google's system: If you are conducting research on machine translation, optical character recognition or other areas where access to a large amount of text is helpful, please contact us. We encourage the use of public domain materials for these purposes and may be able to help.
- + *Maintain attribution* The Google "watermark" you see on each file is essential for informing people about this project and helping them find additional materials through Google Book Search. Please do not remove it.
- + *Keep it legal* Whatever your use, remember that you are responsible for ensuring that what you are doing is legal. Do not assume that just because we believe a book is in the public domain for users in the United States, that the work is also in the public domain for users in other countries. Whether a book is still in copyright varies from country to country, and we can't offer guidance on whether any specific use of any specific book is allowed. Please do not assume that a book's appearance in Google Book Search means it can be used in any manner anywhere in the world. Copyright infringement liability can be quite severe.

About Google Book Search

Google's mission is to organize the world's information and to make it universally accessible and useful. Google Book Search helps readers discover the world's books while helping authors and publishers reach new audiences. You can search through the full text of this book on the web at <http://books.google.com/>

Per
MS
N-3



PROCEEDINGS
OF THE
NATIONAL ACADEMY
OF SCIENCES

OF THE
UNITED STATES OF AMERICA

VOLUME 8, 1922



EXECUTIVE COMMITTEE

EDWIN BIDWELL WILSON, *Managing Editor*

RAYMOND PEARL

VERNON KELLOGG

PUBLISHED MONTHLY

Publication Office: Eschenbach Printing Company, Easton, Pa.

Editorial Office: Massachusetts Institute of Technology, Cambridge

Home Office of the Academy: Washington, D. C.

34317

3. 2. 3, 11 - 2

CONTENTS

NATIONAL ACADEMY OF SCIENCES, BACHE FUND	16
ERRATA	232
INDEX	369

MATHEMATICS

THE RIEMANN GEOMETRY AND ITS GENERALIZATION.....	
<i>By L. P. Eisenhart and O. Veblen</i>	19
RICCI'S PRINCIPAL DIRECTIONS FOR A RIEMANN SPACE AND THE EINSTEIN THEORY	
<i>By L. P. Eisenhart</i>	24
NOTE ON THE DEFINITION OF A LINEAL FUNCTIONAL.....	26
<i>By C. A. Fischer</i>	
ON THE RELATION OF A CONTINUOUS CURVE TO ITS COMPLEMENTARY DOMAINS IN SPACE OF THREE DIMENSIONS.....	33
<i>By Robert L. Moore</i>	
A SOLUTION OF THE LINEAR MATRIX EQUATION BY DOUBLE MULTIPLICATION....	
<i>By F. L. Hitchcock</i>	78
ON THE LOCATION OF THE ROOTS OF THE DERIVATIVE OF A POLYNOMIAL.....	
<i>By J. L. Walsh</i>	139
NORMAL COÖRDINATES FOR THE GEOMETRY OF PATHS.....	192
<i>By O. Veblen</i>	
PRINCIPAL DIRECTIONS IN A RIEMANNIAN SPACE.....	198
<i>By J. L. Synge</i>	
PRINCIPAL DIRECTIONS IN THE EINSTEIN SOLAR FIELD.....	204
<i>By J. L. Synge</i>	
FIELDS OF PARALLEL VECTORS IN THE GEOMETRY OF PATHS....	207
<i>By L. P. Eisenhart</i>	
SPACES WITH CORRESPONDING PATHS.....	233
<i>By L. P. Eisenhart</i>	
NUMBER OF SUBSTITUTIONS OMITTING AT LEAST ONE LETTER IN A TRANSITIVE GROUP.....	238
<i>By G. A. Miller</i>	
THE MEANING OF ROTATION IN THE SPECIAL THEORY OF RELATIVITY.....	
<i>By Philip Franklin</i>	265
NEW PROPERTIES OF ALL REAL FUNCTIONS.....	283
<i>By Henry Blumberg</i>	
GENERALIZED LIMITS IN GENERAL ANALYSIS.....	288
<i>By Charles N. Moore</i>	
PROJECTIVE AND AFFINE GEOMETRY OF PATHS.....	347
<i>By O. Veblen</i>	

ASTRONOMY

THE EFFECTIVE TEMPERATURE OF 16 STARS AS ESTIMATED FROM THE ENERGY DISTRIBUTION IN THE COMPLETE SPECTRUM.....	49
<i>By W. W. Coblentz</i>	
NOTE ON THE PROBLEM OF GREAT STELLAR DISTANCES.....	69
<i>By H. Shapley</i>	
DARK NEBULAE.....	115
<i>By H. N. Russell</i>	
ON THE DISTRIBUTION OF THE VELOCITIES OF STARS OF LATE TYPES OF SPECTRUM.....	141
<i>By G. Strömberg</i>	
INVISIBLE SUN-SPOTS.....	168
<i>By G. E. Hale</i>	
COMPARISON OF THEORY WITH OBSERVATION FOR THE MINOR PLANETS 10 HYGIEA AND 175 ANDROMACHE WITH RESPECT TO PERTURBATIONS BY JUPITER.....	
<i>By A. O. Leuschner</i>	170
A METHOD OF DERIVING THE DISTANCE OF THE A-TYPE STARS.....	
<i>By W. S. Adams and A. H. Joy</i>	173
THE LARGER RESULTS OF 20 YEARS OF SOLAR RADIATION OBSERVATIONS.....	
<i>By C. G. Abbot, F. E. Fowle and L. B. Aldrich</i>	177

STATISTICS

THE VITAL INDEX OF THE POPULATION OF ENGLAND AND WALES, 1838-1920.....	
<i>By R. Pearl and M. H. Burger</i>	71
SEASONAL FLUCTUATIONS OF THE VITAL INDEX OF A POPULATION..	76
<i>By R. Pearl</i>	

THE PROBABLE ERROR OF THE VITAL INDEX OF A POPULATION..	<i>By J. R. Miner</i>	106
ON THE MEASUREMENT OF A PHYSICAL QUANTITY WHOSE MAGNITUDE IS INFLUENCED BY PRIMARY CAUSES BEYOND THE CONTROL OF THE OBSERVER AND ON THE METHOD OF DETERMINING THE RELATION BETWEEN TWO SUCH QUANTITIES.....	<i>By W. A. Shewhart</i>	248
A FURTHER NOTE ON THE AGE INDEX OF A POPULATION.....	<i>By R. Pearl and T. J. LeBlanc</i>	300
THE STABILITY OF THE NORMAL AGE DISTRIBUTION.....	<i>By Alfred J. Lotka</i>	339
A FURTHER NOTE ON THE MATHEMATICAL THEORY OF POPULATION GROWTH....	<i>By Raymond Pearl and Lowell J. Reed</i>	365
PHYSICS		
THE FREQUENCY-SENSITIVITY OF NORMAL EARS..	<i>By H. Fletcher and R. L. Wegel</i>	5
EXPERIMENTS WITH THE VACUUM GRAVITATION NEEDLE.....	<i>By C. Barus</i>	13
ABSORPTION OF SHORT X-RAYS BY ALUMINIUM AND COPPER.....	<i>By W. Duane and K. C. Mazumder</i>	45
THE GRATING SPACE OF MICA AND THE INTENSITIES OF THE SPECTRAL ORDERS....	<i>By Bergen Davis and H. M. Terrill</i>	59
IONIZATION AND RADIATION POTENTIALS AND THE SIZE OF THE ATOM.....	<i>By Bergen Davis</i>	61
PERIODS AND LOGARITHMIC DECREMENT OF THE GRAVITATION NEEDLE UNDER HIGH EXHAUSTION.....	<i>By C. Barus</i>	63
PLANE REFLECTION OF SOUND, AS EXHIBITED BY THE PINHOLE RESONATOR.....	<i>By C. Barus</i>	66
NOTE ON X-RAY SPECTRA.....	<i>By W. Duane and R. A. Patterson</i>	85
A NEW METHOD OF USING X-RAYS IN CRYSTAL ANALYSIS.....	<i>By G. L. Clark and W. Duane</i>	90
ACOUSTIC TOPOGRAPHY VARYING WITH THE POSITION OF THE ORGAN PIPE.....	<i>By C. Barus</i>	96
ON STEERING AN AUTOMOBILE AROUND A CORNER.....	<i>By A. G. Webster</i>	100
CYLINDRICAL DISTRIBUTION OF NODAL STRENGTH AROUND THE PIPE NORMAL.....	<i>By C. Barus</i>	119
IONIZATION AND ABSORPTION EFFECTS IN THE ELECTRIC FURNACE..	<i>By A. S. King</i>	123
THE PHYSICAL EXAMINATION OF HEARING AND BINAURAL AIDS FOR THE DEAF....	<i>By R. L. Wegel</i>	155
THE PRODUCTION OF AN E. M. F. ON CLOSED CIRCUIT BY A LIGHT EFFECT ON ARGENTITE.....	<i>By H. H. Sheldon and P. H. Geiger</i>	161
POSITIVE AND NEGATIVE PINHOLE RESONATORS.....	<i>By C. Barus</i>	163
THE EVALUATION OF QUANTUM INTEGRALS.....	<i>By P. S. Epstein</i>	166
THE SPECTRAL ENERGY DISTRIBUTION AND OPACITY OF WIRE EXPLOSION VAPORS	<i>By J. A. Anderson</i>	231
ELECTRONIC BOMBARDMENT OF NICKEL.....	<i>By H. E. Farnsworth</i>	251
THE CRYSTAL STRUCTURE OF POTASSIUM.....	<i>By L. W. McKeehan</i>	254
PRELIMINARY REPORT ON THE IONIZATION OF POTASSIUM VAPOR BY LIGHT.....	<i>By R. C. Williamson</i>	255
THE ABSORPTION OF LIGHT BY SODIUM AND POTASSIUM VAPORS..	<i>By G. R. Harrison</i>	260
THE ENERGY LOSSES ACCOMPANYING IONIZATION AND RESONANCE IN MERCURY VAPOR.....	<i>By John A. Eldridge</i>	268
THE CRYSTAL STRUCTURE OF BERYLLIUM AND OF BERYLLIUM OXIDE.....	<i>By L. W. McKeehan</i>	270
SOME CASES OF NERVE-DEAFNESS AND THEIR BEARING ON RESONANCE THEORIES OF AUDITION.....	<i>By John P. Minton</i>	274

CONTENTS

v

AN ELECTRON THEORY OF ELECTRIC CONDUCTION IN METALS.....	<i>By Edwin H. Hall</i>	307
STATIC DEFLECTION, LOGARITHMIC DECREMENT AND FIRST SEMI-PERIOD OF THE VACUUM GRAVITATION NEEDLE.....	<i>By C. Barus</i>	313
ON A COMPARISON OF THE RELATIVE SENSITIVENESS OF TELEPHONES.....	<i>By Carl Barus</i>	326
THE EQUILIBRIUM POSITIONS OF THE VACUUM GRAVITATION NEEDLE IN 1921 AND 1922.....	<i>By Carl Barus</i>	327
FURTHER MEASUREMENTS OF STELLAR TEMPERATURES AND PLANETARY RADIA- TION.....	<i>By W. W. Coblentz</i>	330
PROOF OF A THEOREM DUE TO HEAVISIDE.....	<i>By H. A. Lorentz</i>	333
CHARACTERISTICS OF A SHORT WAVE OSCILLATOR AT VERY LOW PRESSURE.....	<i>By L. L. Nettleton</i>	353
THE REFRACTION OF X-RAYS IN CALCITE.....	<i>By Bergen Davis and H. M. Terrill</i>	357
THE COMPRESSIBILITY OF METALS AT HIGH PRESSURES.....	<i>By P. W. Bridgman</i>	361

CHEMISTRY

THE RELATION OF CHALCEDONY TO THE OTHER FORMS OF SILICA.....	<i>By E. W. Washburn and L. Navias</i>	1
THE ACTION OF DIAZOMETHANE ON URACIL.....	<i>By T. B. Johnson, A. J. Hill and F. H. Case</i>	44
THE OXIDATION OF HYDROQUINOL IN THE PRESENCE OF ALIPHATIC AMINES.....	<i>By Rolla N. Harger</i>	57
THE ISOLATION OF PYRIMIDINES FROM THE NUCLEIC ACID OF TUBERCLE BACILLUS	<i>By T. B. Johnson and E. B. Brown</i>	187
THE THERMAL IONIZATION OF GASEOUS ELEMENTS AT HIGH TEMPERATURES: A CONFIRMATION OF THE SAHA THEORY.....	<i>By Arthur A. Noyes and H. A. Wilson</i>	303

GEOLOGY AND PALEONTOLOGY

CORAL REEFS OF THE LOUISIADÉ ARCHIPELAGO.....	<i>By W. M. Davis</i>	7
ADDITIONAL STUDIES IN ARRESTED EVOLUTION.....	<i>By R. Ruedemann</i>	54
NEW FORMS OF LIFE FROM THE SILURIAN.....	<i>By R. Ruedemann</i>	55
THE AVERAGE CHEMICAL COMPOSITION OF IGNEOUS ROCKS.....	<i>By F. W. Clarke and H. S. Washington</i>	108
HESPEROPITHECUS, THE FIRST ANTHROPOID PRIMATE FOUND IN AMERICA.....	<i>By H. F. Osborn</i>	245
RECENT DISCOVERIES ON THE ANTIQUITY OF MAN.....	<i>By H. F. Osborn and C. A. Reeds</i>	246
THE JADES OF MIDDLE AMERICA.....	<i>By Henry S. Washington</i>	319

ZOOLOGY

SKIN TRANSPLANTATION IN FROG TADPOLES.....	<i>By W. H. Cole</i>	29
THE EFFECT OF ETHER UPON THE MIGRATION OF THE SCALE PIGMENT AND THE RETINAL PIGMENT IN THE FISH, FUNDULUS HETEROCLITUS..	<i>By L. C. Wyman</i>	128
THE DIVERSE EFFECTS OF ADRENALIN UPON THE MIGRATION OF THE SCALE PIG- MENT AND THE RETINAL PIGMENT IN THE FISH, FUNDULUS HETEROCLITUS, LINN.....	<i>By A. S. Gilson, Jr.</i>	130
A SIDE LINE IN THE IMPORTATION OF INSECT PARASITES OF INJURIOUS INSECTS FROM ONE COUNTRY TO ANOTHER.....	<i>By L. O. Howard</i>	133
DEGENERATION AND REGENERATION OF THE LATERAL-LINE ORGANS IN AMIURUS NEBULOSUS.....	<i>By M. H. Chambers</i>	186

CONTENTS

	Page
PATHOLOGY.—BACTERICIDAL PROPERTIES OF THE PRODUCTS OF RADIUM EMANATION	317
<i>By John Albert Marshall</i>	
PETROLOGY.—THE JADES OF MIDDLE AMERICA	319
<i>By Henry S. Washington</i>	
PHYSICS.—ON A COMPARISON OF THE RELATIVE SENSITIVENESS OF TELEPHONES	326
<i>By Carl Barus</i>	
PHYSICS.—THE EQUILIBRIUM POSITIONS OF THE VACUUM GRAVITATION NEEDLE IN 1921 AND 1922	327
<i>By Carl Barus</i>	
PHYSICS.—FURTHER MEASUREMENTS OF STELLAR TEMPERATURES AND PLANETARY RADIATION	330
<i>By W. W. Coblentz</i>	
PHYSICS.—PROOF OF A THEOREM DUE TO HEAVISIDE	333
<i>By H. A. Lorentz</i>	
STATISTICS.—THE STABILITY OF THE NORMAL AGE DISTRIBUTION	339
<i>By Alfred J. Lotka</i>	

VOLUME 8

JANUARY, 1922

NUMBER 1

PROCEEDINGS
OF THE
**NATIONAL ACADEMY
OF SCIENCES**

OF THE
UNITED STATES OF AMERICA



EXECUTIVE COMMITTEE

EDWIN BIDWELL WILSON, *Managing Editor*

RAYMOND PEARL

VERNON KELLOGG

PUBLISHED MONTHLY

Publication Office: Eschenbach Printing Company, Easton, Pa.

Editorial Office: Massachusetts Institute of Technology, Cambridge

Home Office of the Academy: Washington, D. C.

Entered as second-class matter June 21, 1920, at the postoffice at Easton, Pennsylvania, under the act of August 24, 1912.
Acceptance for mailing at special rate of postage provided for in section 1103, Act of October 3, 1917. Authorized June 15

Price per annum, \$5.00

EDITORIAL BOARD

The HOME SECRETARY and the FOREIGN SECRETARY of the ACADEMY
The CHAIRMAN and the PERMANENT SECRETARY of the NATIONAL
RESEARCH COUNCIL

J. M. CLARKE
A. L. DAY
R. DODGE
W. DUANE
G. DUNN
N. M. FENNEMAN

A. D. FLINN
F. P. GAY
R. G. HARRISON
L. J. HENDERSON
F. R. LILLIE

E. H. MOORE
W. J. V. OSTERHOUT
F. SCHLESINGER
E. W. WASHBURN
W. M. WHEELER

INFORMATION TO SUBSCRIBERS

SUBSCRIPTIONS TO THE PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES at the rate of \$5.00 per annum should be made payable to the National Academy of Sciences, and sent either to Easton, Pa., or to C. G. ABBOT, Home Secretary, National Academy of Sciences, Smithsonian Institution, Washington, D. C. Single numbers, \$0.50.

PAST publications of the NATIONAL ACADEMY OF SCIENCES are listed in these PROCEEDINGS, Volume III, pp. 743-753, December, 1917. In good part these publications are no longer available for distribution. Inquiries with regard to them should be addressed to the Home Secretary, National Academy of Sciences, Smithsonian Institution, Washington, D. C.

PAST volumes of the PROCEEDINGS may be obtained at five dollars per volume unless the sale of the volume would break a complete set of Volumes I to V. Single numbers may be obtained for fifty cents except where the sale of such numbers would break up a complete volume. Only two hundred complete sets of the Proceedings are available for sale—Volumes I to VI, price \$30.00. Orders should be sent to the Home Secretary, National Academy of Sciences, Smithsonian Institution, Washington, D. C.

The following publications are issued by the NATIONAL RESEARCH COUNCIL. Orders and inquiries should be addressed: Publication Office, National Research Council, 1701 Massachusetts, Ave., Washington, D. C.

(A) THE BULLETIN OF THE NATIONAL RESEARCH COUNCIL, published at irregular intervals; Price \$5.00 per volume of about 500 pages; individual numbers priced variously.

(B) THE REPRINT AND CIRCULAR SERIES OF THE NATIONAL RESEARCH COUNCIL, individual numbers, variously priced.

PROCEEDINGS OF THE NATIONAL ACADEMY OF SCIENCES

Volume 8

DECEMBER 15, 1922

Number 12

PROJECTIVE AND AFFINE GEOMETRY OF PATHS

BY O. VEBLEN

DEPARTMENT OF MATHEMATICS, PRINCETON UNIVERSITY

Communicated, October 31, 1922

1. A path is defined as any curve given parametrically by a set of solutions of the differential equations

$$\frac{d^2x^i}{ds^2} + \Gamma_{\alpha\beta}^i \frac{dx^\alpha}{ds} \frac{dx^\beta}{ds} = 0, \quad \Gamma_{\alpha\beta}^i = \Gamma_{\beta\alpha}^i \quad (1.1)$$

The same system of paths may, however, be defined in terms of the same system of coördinates by another set of differential equations

$$\frac{d^2x^i}{dt^2} + \Lambda_{\alpha\beta}^i \frac{dx^\alpha}{dt} \frac{dx^\beta}{dt} = 0, \quad \Lambda_{\alpha\beta}^i = \Lambda_{\beta\alpha}^i \quad (1.2)$$

The functions $\Gamma_{\alpha\beta}^i$ give rise to one definition of covariant differentiation and infinitesimal parallelism, and the functions $\Lambda_{\alpha\beta}^i$ to a different definition of these operations and relations. But both types of parallelism and of covariant differentiation refer to the same system of paths.

The theorems which state general properties of a system of paths, without restriction as to the scope of the theorems, constitute a projective geometry of paths. The theorems which state properties of the paths and of a particular set of functions, $\Gamma_{\alpha\beta}^i$, i.e., of a particular definition of infinitesimal parallelism, constitute an affine geometry of paths.

The discovery that there can be more than one affine geometry for a given system of paths is due to H. Weyl who also obtained an important tensor which he calls the *projective curvature*. Weyl's results are published in the *Göttinger Nachrichten* for 1921 (p. 99), which has only recently reached this country.

The question whether there could be two sets of differential equations such as (1.1) and (1.2) for the same set of paths was raised by Professor Eisenhart (cf. this volume, p. 233) and answered in the negative under the

assumption that "the parameter s is the same for all the paths." The following paragraphs will serve to show wherein this assumption restricts the problem and also provide a second method of approaching the projective geometry of paths.

2. Let us first see how the differential equations (1.1) and (1.2) can give rise to the same system of paths. Any solution of the differential equations (1.1) can be written in the form (cf. p. 193 of this volume)

$$x^i = q^i + \psi^i(\xi^1 s, \xi^2 s, \dots, \xi^n s) \quad (2.1)$$

in which q^1, q^2, \dots, q^n are the coördinates of an arbitrary point and $\xi^1, \xi^2, \dots, \xi^n$ an arbitrary set of values of $dx^1/ds, dx^2/ds, \dots, dx^n/ds$ at this point. The solution of (1.2) representing the same paths as (2.1) may be written in the form

$$x^i = q^i + X^i(\eta^1 t, \eta^2 t, \dots, \eta^n t) \quad (2.2)$$

in which $\eta^1, \eta^2, \dots, \eta^n$ is a set of values of $dx^1/dt, dx^2/dt, \dots, dx^n/dt$ at the point q . By setting up a correspondence between the values of s and t which correspond to the same point of this path we define s as a function of t . This function depends on the point q and the direction of the path through q . Hence if $y^i = \eta^i t$, we may write

$$s = f(q^1, q^2, \dots, q^n, y^1, y^2, \dots, y^n) \quad (2.3)$$

for the value of s which corresponds to the same point as q^1, q^2, \dots, q^n and y^1, y^2, \dots, y^n . As a transformation of the differential equation this may also be written.

$$s = f(x^1, x^2, \dots, x^n, \frac{dx^1}{dt} t, \frac{dx^2}{dt} t, \dots, \frac{dx^n}{dt} t) \quad (2.4)$$

Let us now multiply (1.1) by $(\partial f / \partial t)^2$ and add a term to each side so as to obtain

$$\frac{d^2 x^i}{ds^2} \left(\frac{\partial f}{\partial t} \right)^2 + \frac{dx^i}{ds} \frac{\partial^2 f}{\partial t^2} + \Gamma_{\alpha\beta}^i \frac{dx^\alpha}{ds} \frac{dx^\beta}{ds} \left(\frac{\partial f}{\partial t} \right)^2 = \frac{dx^i}{ds} \frac{\partial^2 f}{\partial t^2} \quad (2.5)$$

which is the same as

$$\frac{d^2 x^i}{dt^2} + \Gamma_{\alpha\beta}^i \frac{dx^\alpha}{dt} \frac{dx^\beta}{dt} = \frac{dx^i}{dt} \Theta(x^1, x^2, \dots, x^n, \frac{dx^1}{dt}, \frac{dx^2}{dt}, \dots, \frac{dx^n}{dt}, t) \quad (2.6)$$

a differential equation in which t has exactly the same significance as in (1.2).

3. Let us now subtract (1.2) from (2.6). The result is

$$\varphi_{\alpha\beta}^i \frac{dx^\alpha}{dt} \frac{dx^\beta}{dt} = \Theta \frac{dx^i}{dt} \quad (3.1)$$

in which we are defining the functions $\varphi_{\alpha\beta}^i$ by the equations

$$\varphi_{\alpha\beta}^i = \Gamma_{\alpha\beta}^i - \Lambda_{\alpha\beta}^i \quad (3.2)$$

The functions thus defined are the components of a tensor which is contravariant with respect to i and covariant with respect to α and β . This is because a transformation to new independent variables z^1, z^2, \dots, z^n , changes the Γ 's according to the formula

$$\Gamma_{\alpha\beta}^i(z) = \Gamma_{\gamma\delta}^j(x) \frac{\partial x^\gamma}{\partial z^\alpha} \frac{\partial x^\delta}{\partial z^\beta} \frac{\partial z^i}{\partial x^j} + \frac{\partial^2 x^i}{\partial z^\alpha \partial z^\beta} \frac{\partial z^j}{\partial x^j} \quad (3.3)$$

This tensor defines a covariant vector by means of the formula

$$\varphi_\alpha = \frac{1}{n+1} \varphi_{i\alpha}^i \quad (3.4)$$

By eliminating Θ from the equations (3.1) we obtain

$$\left(\varphi_{\alpha\beta}^i \frac{dx^i}{dt} - \varphi_{\alpha\beta}^j \frac{dx^j}{dt} \right) \frac{dx^\alpha}{dt} \frac{dx^\beta}{dt} = 0 \quad (3.5)$$

Remembering that the values of $dx^1/dt, dx^2/dt, \dots, dx^n/dt$ are arbitrary it is easy to infer from (3.5) that

$$\varphi_{\alpha\beta}^i = \delta_\alpha^i \varphi_\beta + \delta_\beta^i \varphi_\alpha \quad (3.6)$$

where δ_α^i is 1 or 0 according as $i = \alpha$ or $i \neq \alpha$.

When (3.6) is substituted in (3.1) there results

$$2 \varphi_\gamma \frac{dx^\gamma}{dt} = \Theta. \quad (3.7)$$

4. Conversely it can be shown that if we start with the differential equation (1.1) and any covariant vector φ_α , we can find a second differential equation of the type (1.2) which defines the same set of paths as (1.1). The first step is to define the functions $\varphi_{\alpha\beta}^i$ by means of equations (3.6). Since δ_α^i is a tensor and φ_α is a covariant vector it follows that $\varphi_{\alpha\beta}^i$ is a tensor of the third order. We then define $\Lambda_{\alpha\beta}^i$ by means of the equations (3.2) and write the differential equations (1.2) which are by definition identical with

$$\frac{d^2 x^i}{dt^2} + \Gamma_{\alpha\beta}^i \frac{dx^\alpha}{dt} \frac{dx^\beta}{dt} = 2 \varphi_\alpha \frac{dx^\alpha}{dt} \frac{dx^i}{dt} \quad (4.1)$$

If we compare (4.1) with (2.5) we see that (4.1) is obtainable from (1.1) by means of the substitution

$$s = e^{2\varphi_\gamma \frac{dx^\gamma}{dt}} + h(x) \quad (4.2)$$

where $h(x)$ is an arbitrary function of x^1, x^2, \dots, x^n .

It has been proved that: (1) If (1.1) and (1.2) are differential equations which define the same system of paths then the functions $\Gamma_{\alpha\beta}^i$ and $\Lambda_{\alpha\beta}^i$ are related by the equations

$$\Gamma_{\alpha\beta}^i - \Lambda_{\alpha\beta}^i = \varphi_{\alpha\beta}^i = \delta_{\alpha}^i \varphi_{\beta} + \delta_{\beta}^i \varphi_{\alpha} \quad (4.3)$$

in which φ_{α} is a vector. (2) If (1.1) are the differential equations of a system of paths and φ_{α} is any covariant vector then there exists a second set of differential equations (1.2) for the same system of paths, the functions $\Lambda_{\alpha\beta}^i$ being defined by (4.3).

The formula (3.7) reduces to that obtained by Eisenhart on page 235 of this volume in the case when s is a function of t alone. This makes it evident that his work must be taken as referring to a geometry in which there is an element of distance ds defined by means of functions of position. In this sense his theorem on page 235 will be found not to be in contradiction with the theorem (2) above.

A DIRECT METHOD OF TESTING COLOR VISION IN LOWER ANIMALS

BY W. F. HAMILTON

DEPARTMENT OF PHYSIOLOGY, YALE UNIVERSITY

Communicated, October 26, 1922

1. When spectral lights of different wave-lengths are directed on the photoreceptors of lower organisms, their reactions show that one of the two lights is, physiologically, the more effective. Even when the unequal distribution of energy in the spectrum is equated for, there still remains a difference in the physiological effectiveness of the stimuli. (Laurens and Hooker.¹) This difference must be dependent upon wave-length. But it is also evident to the color blind eye (Koenig²). We cannot therefore argue that the differential response on the part of the organism is proof of color division.

Researches which have approached the question of color vision in lower organisms by means of *conditioned behavior* have not been done with pure spectral lights. Even when pure spectral lights are used, the difference in brightness must be equalized, not, however, in accordance with the luminosity curve of the human eye, but with that of the organisms whose color vision is to be tested. Obviously it is difficult to obtain a luminosity curve free from extraneous factors from the reactions of organisms sufficiently complex and intelligent to exhibit conditioned behavior.

The method of conditioned behavior is not applicable to lower organisms and it meets with difficulties when applied to higher animals. Therefore, if we would study the evolution of reactions to color, we must cast about for another method.

Most theories of color vision postulate receptive systems maximally sensitive to certain wave-lengths which differ in some manner from those which are maximally sensitive to other wave-lengths. The difference, according to some theories, lies in the receptive substance of the eye (Young-Helmholtz, Hering); according to others, there are different "color perceiving" mechanisms in the brain which serve for perception of the different fundamental colors. (Edridge Green,³ Houstoun,⁴ and the various "Zone Theories.") Work on the human eye (Burch,⁵ Abney,⁶ Frank Allen⁷ and others) shows that these receptive systems can be separately fatigued by strong monochromatic light.

If we can get evidence that the photo receptors of lower organisms can be differentially fatigued in a similar manner, it would indicate that the receptive system which is maximally sensitive to certain wave-lengths differs in some respect from the receptive system which is maximally sensitive to other wave-lengths, and this could be construed into something analogous to color vision. The following is a preliminary report of the results of experiments recently performed with this in view.

2. The animal decided upon was the fruit-fly, *Drosophila*. Specimens were kept in the dark over night and used in the dark adapted state. The apparatus consisted of two Hilger wave-length spectrometers** used as monochromatic illuminators and so set as to throw beams of light into opposite ends of a small horizontal tubular glass cell containing the flies. The sources of light were two 500-watt Mazda C stereoptican lamps.

The intensities of the light beams, differing as to wave-length were so regulated that the animals were not oriented in either direction. The lights were physiologically equal. As soon as this was accomplished one beam of light, called for convenience the "stimulating light," was intercepted by a screen and the animals subjected to the action of the other ("fatiguing") light. As soon as a marked decrease in the positive reaction of the flies to the "fatiguing" light was observed, the screen was removed and the animals subjected to the "stimulating" light again.

As soon as this was done, the animals usually oriented and crawled toward the "stimulating" light in spite of the continued action of the "fatiguing" light. The operation of the "fatiguing" light had thus upset the physiological balance between the two lights by fatiguing the apparatus maximally sensitive to this light, more than the apparatus maximally sensitive to the "stimulating" light. When such a test had been complete, it was repeated immediately, with the light that had been used as

the "fatiguing" light arranged to "stimulate" and vice versa. In the second test, the animals were, of course, not dark adapted.

If, however, the two lights were of the same or nearly the same wave-lengths the results were quite different. The lights were balanced, the "stimulating" light intercepted, and the animals subjected to the "fatiguing" light. There was evidence of fatigue, but when the "stimulating" light was again allowed to fall on the flies, it was easily seen that the fatigue was not differential. The animals distributed themselves at random, between the two ends, just as they did when the lights were first balanced.

The spectrum was then scaled. The smallest difference in wave-length which showed conclusive evidence of differential fatigue was the unit used. Beginning at 385 $m\mu$, which is as low as the calibration of the instruments extend, the values were found as follows:

385-410, 410-430, 430-450, 450-500 $m\mu$

The last determination was rather doubtful. At the intensity available the orienting effect of light above 500 $m\mu$ was insufficient to give conclusive evidence of differential fatigue. The distribution of these points and the fact that "hue perception" is at its maximum between 410 $m\mu$ and 450 $m\mu$ makes it appear that there may be two receptor systems, one for the blue-violet and one for the blue-green. Evidence for a third component seems to be lacking. According to Steindler⁸ the hue perception curve of the normal human eye indicates the existence of three receptor systems.

3. It is planned to extend this work to other organisms and to work out the energy-time relationships of differential fatigue with different pairs of wave-lengths at different intensities.

The method should prove of value in analyzing the wave-length receptors of lower organisms, and its modifications and applications are numerous. It is applicable to the investigation of the pupil reactions, of the action current of the optic nerve and to similar problems. Perhaps one of its most interesting applications would be to ascertain the course of evolution of the color sense among lower organisms by means of photic orientation in crossed beams of light. The chief value of the method is its objectiveness.

¹ Laurens, H., and Hooker, H. O., 1920, Studies on the relative physiological value of spectral lights. *J. Exper. Zool.*, 30, 345-68.

² Koenig, *Ges. Abhandlungen*, Berlin.

³ Edrige Green, F. W., 1920, *The Physiology of Vision*. London.

⁴ Houstoun, R. A., 1916, A theory of color vision. *Proc. Roy. Soc. London*, A92, 424-32.

⁵ Burch, G. T., 1899, On artificial temporary color blindness with an examination of the color sensation of 109 persons. *Proc. Roy. Soc.*, B191, 1-34.

⁶ Abney, Sir W. de W., 1913, *Researches in Color Vision*. London.

¹ Allen, Frank, 1920, Persistence of vision and the primary color sensations. *Amer. J. Physiol. Optics*, 1, 94-134.

² Steindler, 1906, Der Farbenempfindlichkeit des normalen und farbenblinden Auges. *Sitz. Wiener Akad.*, 105, 11a, 115-6.

^{**} One of these instruments belongs to this Laboratory, while the other was loaned to us by the kindness of Professor G. H. Parker, Harvard University.

CHARACTERISTICS OF A SHORT WAVE OSCILLATOR AT VERY LOW PRESSURES

BY L. L. NETTLETON

DEPARTMENT OF PHYSICS, UNIVERSITY OF WISCONSIN

Communicated October 11, 1922

Various investigators have experimented with three element vacuum tubes oscillating at a wave-length of the order of one meter. All of them have used commercial tubes, having a cylindrical arrangement of filament, grid, and plate.

Barkhausen and Kurz¹ discovered a type of oscillation which was apparently due to the motion of electrons in the tube itself, and was independent of the inductance and capacity of the external circuit. The computed time for an electron to travel across the tube, from filament to plate and back, under the potentials applied, agreed fairly well with the period of the waves as measured by means of Lecher wires coupled to the tube.

Whiddington² experimented with a similar type of oscillation, but he observed much lower frequencies and attributed the observed effects to the motion of ions instead of electrons. An explanation was given which involved a discontinuous emission from the filament.

Gill and Morrell³ have experimented extensively with the Barkhausen type of oscillation, using commercial tubes made by the Marconi company, and have given an explanation of the effect which involves a natural mode of oscillation of the electrical system connected to the tube.

The work of Barkhausen and Kurz and that of Gill and Morrell was done with so-called "hard" tubes, which in the process of manufacture have been thoroughly heated to remove occluded gas and then sealed off from the vacuum pump. These men have assumed that they were dealing with a purely electronic phenomenon, and that no gaseous ionization was concerned.

The essential departure in the experimental arrangement in the present investigation was the use of a tube which was left permanently connected to the vacuum pumps, and was made with a ground joint to make the in-

side parts accessible for modification, replacement of burned out filaments, etc. Also, more power was used than by other investigators, electron currents as high as 300 milamperes and at voltages up to 700 being used at times.

The electrical connections were made as indicated in figure 1, except that the filament heating circuit is not shown. The Lecher wires, L, L , were closely coupled to, but not in metallic contact with, the grid and plate leads, which were brought straight in through the sides of the tube. A sliding bridge, B , across the Lecher wires, carried a crossed wire thermocouple which was connected to a sensitive galvanometer by long, twisted leads. This thermocouple served to measure the strength of oscillations, and by




FIGURE 1

sliding the bridge along the wires, the points of maximum deflection gave the loops of the standing waves, and the distance between these points gave the half wave-length of the oscillations. In the discussion which follows, by strength of oscillations is meant the deflection of the thermocouple galvanometer when the bridge position is adjusted so that this deflection is a maximum.

Oscillations of the Barkhausen type, at wave-lengths of from 50 to 200 cm., were obtained. The unusual characteristic of a tube when oscillating in this way is that there is a negative current to the plate, which

may flow against a negative potential of 100 volts or more with respect to the filament, the grid being at a positive potential of several hundred volts. If the plate is simply connected to the filament through a voltmeter, this voltmeter may read as high as 150 volts, or if connected through an ammeter, the negative current to the plate may be as much as 25% of the total electron current from the filament. The magnitude of this negative current or voltage varies greatly with the temperature of the filament, grid voltage, and gas pressure. There is never oscillation in the Lecher wires without a negative current or potential on the plate, but this negative current may be present and yet the Lecher wires show very feeble or no oscillations. The characteristics of this negative current and the associated oscillations have been studied as affected by changes of grid voltage, of electron current, and of gas pressure.

It was found, very unexpectedly, that both the negative plate current and any sign of oscillation, as indicated by the Lecher wires, ceased abruptly




FIGURE 2

at very low pressures. This is shown in the curves of figure 2 for plate voltages, and in figure 3 for the strength of oscillation in the Lecher wires. These observations were all made with the grid at 250 volts and the filament heating currents such that the electron current was 200 milliamperes. The pressure of the residual gas in the tube was measured with an ionization manometer. The pressures of hydrogen and air were controlled by adjusting the flame under the mercury vapor pump, with liquid air on the trap to keep out mercury vapor. The mercury vapor pressure was controlled by keeping the liquid air trap (with a pool of mercury at

the bottom) at different temperatures, the mercury vapor pump operating at its maximum rate to reduce the pressure of any gas to the minimum. The curves are plotted with the readings of the ionization manometer as abscissae. This was done because it is probable that the essential thing is not the presence of the gas itself, but of gaseous ions. Since different gases at the same pressure give different amounts of ionization when under the same ionizing agent, it seemed better to plot their ionization instead of pressure. The pressure corresponding to a given ionization current is




FIGURE 3

for mercury about half, and for hydrogen about twice that for air. The pressure scale for air is given on the curves, so by multiplying and dividing by 2 the approximate pressures of hydrogen and mercury, respectively, will be obtained.

These curves apparently indicate that a small amount of ionization is necessary for this type of oscillation. The kind of gas does not seem important. Since the gas pressure at which the oscillation stops is so low, being less than .00005 mm. there can be no connection with the mean free path of the ions, for at these pressures the path is many times longer than the dimensions of the tube. It is not likely that the commercial tubes used by other investigators are sufficiently exhausted to give any indication of the dropping off of the curves at the low pressure side. So far, no satisfac-

tory explanation has been found for the profound effect of such a small amount of gas or the behavior of the tube.

The investigation is being continued and a more detailed account will be published in the near future.

¹ Barkhausen and Kurz, *Phys. Zs., Leipzig*, **21**, No. 1, Jan. 1920, p. 1.

² Whiddington, *Radio Review*, Nov. 1919, p. 53.

³ Gill and Morrell, *Phil. Mag.*, **44**, No. 259, July 1922, p. 161.

THE REFRACTION OF X-RAYS IN CALCITE

BY BERGEN DAVIS AND H. M. TERRILL

DEPARTMENT OF PHYSICS, COLUMBIA UNIVERSITY, N. Y.

Communicated, October 26, 1922

The effect of refraction in X-ray spectra has been discussed by Stenström,¹ who made some determinations of the index of refraction from the relative displacement of the several orders. In most cases the effect was too small to admit of measurement, but for sugar and gypsum crystals he obtained some values for wave-lengths greater than 2.5 Å.

The present paper applies a modification of the same method to the reflection from calcite of the $K\alpha_1$ line of Molybdenum, 70783 Å.

Since we measure the angle from the crystal face, and not the normal, the customary equation for the index of refraction becomes

$$\nu = \frac{\cos \theta}{\cos \theta'} \quad (1)$$

where θ is the glancing angle outside the crystal and θ' the angle of the beam inside. We will use a subscript to indicate orders higher than the first. Placing $\nu = 1 - \delta$, Stenström computes the values of δ from the equation

$$\delta = \frac{\left(\frac{\sin \theta_m}{m}\right)^2 - \left(\frac{\sin \theta_n}{n}\right)^2}{2\left(\frac{\cos \theta_m}{m}\right)^2 - 2\left(\frac{\cos \theta_n}{n}\right)^2}$$

where m and n are any two orders.

As the shift is small in any case, it seemed desirable to express the value of δ directly in terms of angle, in order to more readily determine the effect of errors of observation which are liable to be of the same order of magnitude as δ itself. Now from (1) we have

$$\sin^2 \theta' = 1 - \nu^{-2} \cos^2 \theta$$

Since δ is small, we write

$$\nu^{-2} = 1 + 2\delta$$

whence

$$\sin \theta = [1 - \cos^2 \theta (1 + 2\delta)]^{1/2} = [\sin^2 \theta - 2\delta \cos^2 \theta]^{1/2}$$

and again neglecting terms containing higher powers of δ .

$$\sin \theta' = \theta - \delta \frac{\cos^2 \theta}{\sin \theta} \quad (2)$$

Taking arc sin of each side, we have since δ is small,

$$\begin{aligned} \theta' &= \theta - \delta \frac{\cos^2 \theta}{\sin \theta} \cdot \frac{1}{\cos \theta} \\ \theta' &= \theta - \delta \cot \theta \end{aligned} \quad (3)$$

Now in this equation, θ is the crystal angle measured, hence by determining the angles of any characteristic line for several orders, it will be possible to determine δ , or at least to find the limits of its magnitude. The angles being measured, those of higher orders are reduced to the equivalent first order angle, so that all angles may be compared directly. The proportion that each is affected by refraction being known from (3), the absolute value of δ is readily determined.

Instead of using (3), it is possible to substitute (2) in the relation

$$n \lambda = 2 d \sin \theta_n'$$

obtaining, since δ is small

$$\frac{n \lambda}{2 \sin \theta_n} = d (1 - \delta \cot^2 \theta_n) \quad (4)$$

thus determining δ from the increase in apparent grating space with order. However, since the limits of error have to be carried through in the computations, the previous method seems to be more convenient.

The spectrometer used has been described elsewhere.² The tube was of the water cooled type with a molybdenum target. Measurements of θ were made for the first three orders of the $K\alpha_1$ line, using a crystal of clear Iceland spar. The measurements were gone over several times and it was found that the angles could be repeated to within 15". This is just about the limit of accuracy of the instrument which was determined by other methods to be about 20".

The following mean values were obtained.

$$\begin{aligned} \theta &= 6^\circ 42' 43'' \pm 10'' \\ \theta_2 &= 13^\circ 30' 45'' \pm 10'' \\ \theta_3 &= 20^\circ 31' 22'' \pm 10'' \end{aligned}$$

where the additional terms represent the limits of error.

It should be remarked that if the crystal is not mounted so that the mean reflecting plane is directly over the center of rotation, and the face of the crystal be slightly curved, a shift of the lines will be obtained, that might be taken for refraction. To eliminate this possibility, the crystal was mounted on a slide with a micrometer screw attached, so that it might be moved in a direction perpendicular to its face. The crystal being mounted as nearly as possible over the center of rotation by mechanical methods, the adjustment was changed by small fractions of a millimeter, reading each time on both sides the crystal and chamber angles of the α_1 line. If curvature is present, the crystal angle will change as the crystal is moved. Twice the crystal angle is plotted against the readings of the micrometer head and on the same sheet, the chamber angle is also plotted against these readings. The intersection of the two lines gives the proper micrometer setting, since the crystal must be over the center, when the chamber angle is exactly twice the crystal angle. With the crystal used, it was found that the curvature was very slight, a shift of less than one second of arc being found within the range of adjustment. This was checked for each order.

We proceed then to reduce the angles measured to the corresponding first order angles. Now $\theta_n = \theta'_n + \Delta$ where Δ includes the slight increase due to refraction and also the error of observation. Since Δ is small

$$\sin \theta_n = \sin \theta'_n + \Delta \cos \theta_n$$

$$\frac{\sin \theta_n}{n} = \sin \theta'_n + \Delta \frac{\cos \theta_n}{n}$$

$$\text{arc sin} \left(\frac{\sin \theta_n}{n} \right) = \theta' + \Delta \frac{\cos \theta_n}{n \cos \theta}$$

$$\theta' = \text{arc sin} \left(\frac{\sin \theta_n}{n} \right) - \Delta \frac{\cos \theta_n}{n \cos \theta}$$

and the coefficient of Δ can be computed for each order, it being .49 for the second and .31 for the third. Multiplying the error limits by these coefficients and introducing the refraction from (3) which is also multiplied by them, we have the following values for θ' :

$$\begin{aligned} \theta' &= 6^\circ 42' 43'' \pm 10'' - 1''.76 \cdot \delta \cdot 10^6 \\ &= 6^\circ 42' 33'' \pm 5'' - 0''.63 \cdot \delta \cdot 10^6 \\ &= 6^\circ 42' 38''.5 \pm 3'' - 0''.27 \cdot \delta \cdot 10^6 \end{aligned}$$

and we have to determine δ from these equations.

Plotting the values of the angles against the order, using lines covering the limits of error instead of points to locate the extremities of the ordinates,

we have to pass through these lines a curve whose ordinates are proportional to the coefficients of δ . Now the values found do not lie well on any curve of this type, the second order being entirely too low. The curves that fitted most nearly gave a value of $\delta = 3 \times 10^{-6}$ while the error limits would permit of values ranging from zero to several times this. It will be seen that this corresponds to a shift of the first order of $5''$, so that for this wave-length, the effect of refraction is very slight.

This value $\delta = 3 \times 10^{-6}$ necessarily contains large possible errors. It is, however, of the same order of magnitude as that obtained by the method of total reflection.

An independent determination of δ was obtained by measuring the angle of total reflection as suggested by the experiments of A. H. Compton³ on total reflection from glass and lead. The ionization chamber was shielded from the direct beam and the spectrum searched at angles close to grazing incidence. Total reflection appeared as a line, approximately $4'$ wide at half maximum, slightly unsymmetrical and of an intensity about 25 times the surrounding values which coincided with the leak. To test if this line consisted principally of the monochromatic radiation corresponding to the $K\alpha$ line, a zirconium screen was introduced. This diminished the intensity without materially changing the shape of the line or shifting its position.

Measurements were made on each side of the zero position, and the mean of the readings gave $6' 30''$, the value being good to $30''$. Now for total reflection, putting $\theta' = 0$ in (1) we have

$$\nu = \cos \theta \text{ or } \delta = \frac{\sin^2 \theta}{2}$$

whence

$$\delta = 1.7 \pm .5 \times 10^{-6}$$

Comparing this with the theoretical value from the Lorentz equation

$$\delta = \frac{n e^2}{2 \pi m} \cdot \frac{1}{\nu}$$

we take the density of calcite as 2.71 giving 81.5×10^{22} electrons per cubic centimeter, and place ν equal to the frequency of the molybdenum $K\alpha$ line, whence $\delta = 1.85 \times 10^{-6}$.

Determinations of the total reflection made with a second crystal gave results which checked with the preceding. Incidentally it developed that the face of this second crystal had been ground at a slight angle to the reflecting planes, the crystal zero determined by the angle of total reflection differing from that obtained from the line spectra by an angle of $44'$. This suggests an extension of the method of determining the refraction

from the shift of the spectral lines, for if the face of the crystal were ground at the proper angle to the reflecting planes, the shift would be increased enough to make accurate determinations possible.

¹ *Experimentelle Untersuchungen der Röntgenspektira*, Lund 1919.

² *J. Amer. Opt. Soc.*, May 1922.

³ *Physic Rev.*, Ithaca, July 1922.

THE COMPRESSIBILITY OF METALS AT HIGH PRESSURES

By P. W. BRIDGMAN

JEFFERSON PHYSICAL LABORATORY, HARVARD UNIVERSITY

Communicated October 17, 1922

This note briefly summarizes results which will be found in full detail in a forthcoming number of the PROCEEDINGS of the American Academy of Arts and Sciences. A considerable part of the expenses of the experiment was defrayed by a generous appropriation from the Rumford Fund of the American Academy.

A new method has been devised by which the difference of the linear compressibility of a solid in the form of a rod or wire and the linear compressibility of iron may be measured with an accuracy high enough to give the variation of compressibility with pressure or temperature. In addition, I have improved the method which I previously used¹ for the measurement of the absolute linear compressibility of iron, so that I have now measured the change of linear compressibility with pressure with some accuracy, and I have also obtained improved values for the change of linear compressibility with temperature. By combining these absolute measurements on iron with the measurements of the difference of compressibility the absolute linear compressibility of a number of metals has been obtained.

If the metal is equally compressible in all directions, the absolute volume compressibility may be calculated from the linear compressibility. The great majority of metals crystallize in the cubic system or in the hexagonal close packed arrangement of spheres, and for these the compressibility is equal in all directions. Table I gives the compressibility, χ (defined as $(\partial v / \partial p)_\tau$ where v is the volume of the amount of the metal that under standard conditions occupies 1 cc.) at atmospheric pressure at 30°, the pressure derivative of the compressibility, and the change of compressibility per degree rise of temperature for a number of such metals. The unit of pressure is the kilogram per square centimeter, the pressure range of the experiments was 12000 kg./cm.², and measure-

TABLE I

SUB- STANCE	$-\chi_{\text{AT } 30^\circ}$ $\times 10^7$	$-\frac{1}{\chi} \left(\frac{\partial \chi}{\partial p} \right)_T$ $\times 10^4$	$+\frac{1}{\chi} \left(\frac{\partial \chi}{\partial T} \right)_P$ $\times 10^4$	SUB- STANCE	$-\chi_{\text{AT } 30^\circ}$ $\times 10^7$	$-\frac{1}{\chi} \left(\frac{\partial \chi}{\partial p} \right)_T$ $\times 10^4$	$+\frac{1}{\chi} \left(\frac{\partial \chi}{\partial T} \right)_P$ $\times 10^4$
Al	13.43	0.74	5.4	Ni	5.29	0.802	2.5
Ca	56.97	1.66	6.0	Pd	5.28	0.80	1.3
Ce	35.74	1.06	0.4	Pt	3.60	1.00	3.7
Co	5.39	0.78	3.0	K	356.5	16.1	35.1
Cu	7.19	0.72	4.6	Ag	9.87	0.90	3.8
Ge	13.78	0.98	-2.3	Na	156.2	3.8	22.
Au	5.77	1.08		Sr	81.87	1.78	2.2
Fe	5.87	0.72	2.3	Ta	4.79	0.10	1.0
Pb	23.73	1.46	5.6	W	2.93	1.02	1.5
Li	86.92	2.24	7.2	Tl*	34.2	3.36	12.2
Mg	29.60	1.36	2.8	Ur†	9.66	0.52	-2.5
Mo	3.47	0.72	0.6				

* The crystal system of Tl has not yet been determined. The compressibility listed above was determined from a single specimen and is very materially higher than the value of other observers. It is therefore probable that the crystal system of Tl is not cubic.

† Crystal system of Ur not yet determined.

ments were made at two temperatures, 30° and 75° . The relation between volume and pressure of all these metals, except Na, K, and Tl, could be represented within the accuracy of the experiments by a second degree equation in the pressure. For the three exceptional metals, however, the relation is more complicated, and the coefficients listed in table I are only the initial values at atmospheric pressure. Of the above list of metals, the compressibility of Ge, Sr, and Ur seems never to have been measured, even at low pressures.

It is to be noticed that the order of magnitude of the relative change of compressibility with pressure or temperature is the same for all the metals. The compressibility of every metal decreases numerically with rising pressure, and in nearly every case increases with rising temperature. There are two apparent exceptions to this last statement in the cases of Ge and Ur. It is possible that the result found for Ur may be due to experimental error, but this explanation seems less likely in the case of Ge. This metal is known to be abnormal in other respects.

If the metal does not crystallize in a system whose compressibility is the same in all directions, then a complete description of the distortion under pressure requires that measurements be made of the linear compressibility in a number of directions of orientations known with respect to the crystal structure. This elaborate program I have not as yet been able to carry through, but I have been able to establish in several cases that there are very considerable differences of compressibility in different directions. These differences are shown by the fact that measurements on different

specimens of the same metal do not yield consistent results, unlike measurements on different specimens of cubic metals. If a non-cubic metal is measured first as a casting and then as an extruded rod, different results will nearly always be found. Results on two different castings, even, are not the same, unless the method of cooling the castings has been the same, so as to ensure the same average orientation of the crystal grains. By measuring the linear compressibility of non-cubic metals prepared in different ways, it is possible to set a lower limit to the differences of compressibility that would be found in different directions in a single crystal. This was the method adopted, and in table II are the results for several non-cubic metals. The values tabulated are the linear compressibility, k (defined as $(\partial l / \partial p)_T$, where l is a length parallel to one of the sides of a body of

TABLE II

SUBSTANCE		$-k \text{ AT } 30^\circ$ $\times 10^7$	$-\frac{1}{k} \left(\frac{\partial k}{\partial p} \right)_T$ $\times 10^4$	$+\frac{1}{k} \left(\frac{\partial k}{\partial T} \right)_p$ $\times 10^4$
Bi	cast	7.67	0.97	0.9
	extruded	11.78	1.82	3.7
Sb	cast	6.80	1.39	1.0
	cast	4.89	0.93	1.7
Cd	cast	4.72	1.15	5.6
	cast	6.51	1.23	7.4
	uni-crystalline	2.86	0.94	
Sn	extruded	6.51	1.12	6.6
	cast	5.67	0.71	4.7
Te	cast	-2.36		
Zn	cast	1.66		
		5.30		
		7.13		

metal which under standard conditions is a cube one centimeter on a side) at 30° at atmospheric pressure, and its pressure and temperature derivatives.

The three measurements on zinc were on pieces cut from the same original single crystal, oriented so as to give three directions mutually at right angles. For zinc, therefore, these numbers give a fair idea of the difference of compressibility in different directions in the same crystal. It is surprising that the difference is so large. The mean of the values listed above for three perpendicular directions is nearly the same as one-third the average cubic compressibility found by other observers, confirming the essential correctness of these measurements.

The negative compressibility of tellurium in a particular direction is perhaps unexpected, and so far as I know this is the first case found of this kind of behavior, although a somewhat analogous instance, the negative Poisson's ratio of pyrites, has been found by Voigt.

A number of theoretical considerations are suggested by these measurements of compressibility. In the first place, and most important, I have shown in the detailed paper by a dimensional argument that the correct order of magnitude for the compressibilities of all these metals may be found by taking over for the metal the same picture of the structure that has given such suggestive results when applied to the salts, namely the picture of the metal as composed of two interpenetrating lattices, one of positively charged atomic ions, and the other of negative electrons, the electron lattice in the case of the metal taking the place of the lattice of negatively charged atoms in the case of the salts, as for example the chlorine lattice in NaCl. The same conclusion as to the structure of a metal has been recently drawn by Kraus² on the basis of a large number of chemical facts.

The expression of Born³ for the compressibility of salts, in which a repulsive potential between atoms as the inverse ninth power is found to give good values for the compressibility, cannot be extended to metals, for neither is the absolute value of the compressibility nor its variation with pressure such as would be given by this sort of a potential. There seems to be no simple repulsive potential that will account for the compressibility data for the various metals. It is possible, however, to represent the repulsive potential by an arbitrary function, and then to find the numerical value of several of its derivatives from experimental data. For those metals which crystallize face-centered cubic, and which therefore probably consist of interpenetrating lattices of ions and electrons of the type of NaCl, it is possible to calculate the grating constants, and to evaluate the first three derivatives of the repulsive potential in terms of the absolute dimensions of the crystal, the compressibility, and the change of compressibility with pressure. These derivatives for a number of metals are listed in the detailed paper. The sign of the derivatives is alternately positive and negative, and their successive absolute values differ by factors of the order of magnitude of the distance of separation of atomic centers. This is all as one would expect. To make more detailed use of these numerical values of the derivatives demands that more attention be paid to the details of the atomic structure than we have hitherto been able to do.

An effect recently discussed by Schottky,⁴ namely a distortion of the atoms under pressure, I have shown by a detailed numerical examination to be of such a magnitude as to essentially modify the simple calculations of compressibility given by Born.

By the use of simple thermodynamics I have been able to modify the arguments of Born, which applied only at the absolute zero of temperature, so as to take account of the effect of temperature on the compressibility to be calculated in terms of the lattice structure. The new terms introduced by temperature are not large enough to seriously modify Born's argument.

Finally, with the new experimental data we are now in a position to calculate for the first time the changes in the specific heat of a metal at high pressures. It appears that the specific heat at constant volume decreases under pressure by an amount of the same order of magnitude as the change of volume under the same pressure, but in most cases by a factor several fold greater.

¹ Bridgman, P. W., *Proc. Amer. Acad., Boston*, **44**, 1909 (255-279), and **47**, 1911 (362-368).

² Kraus, C. A., *J. Amer. Chem. Soc., Easton, Pa.*, **44**, 1922 (1216-1238).

³ Born, M., *Verh. Deut. Phys. Ges., Braunschweig*, **21**, 1919 (533-538).

⁴ Schottky, W., *Phys. Zs., Leipzig*, **21**, 1920 (232-141).

A FURTHER NOTE ON THE MATHEMATICAL THEORY OF POPULATION GROWTH¹

BY RAYMOND PEARL AND LOWELL J. REED

In an earlier paper in these PROCEEDINGS we² showed that the expression

$$y = \frac{be^{ax}}{1 + ce^{ax}} \quad (i)$$

gives an excellent fit to the known population growth of the United States since 1790. Since the first paper was published, we have extended and generalized our ideas on population growth with the results herein presented in skeleton outline. A more extended paper, giving a full discussion of our new results and of the pertinent literature is in press in *Metron*.

Considered generally, the curve

$$y = \frac{b}{e^{-ax} + c}$$

may be written

$$y = \frac{k}{1 + me^{ka'x}}, \quad (ii)$$

where

$$k = b/c, \quad m = 1/c, \quad \text{and} \quad ka' = -a.$$

Now the rate of change of y with respect to x is given by

$$\frac{dy}{dx} = -a'y (k-y)$$

or

$$\frac{\frac{dy}{dx}}{y(k-y)} = -a'. \quad (\text{iii})$$

If y be the variable changing with time x (in our case population) equation (iii) amounts to the assumption that the time rate of change of y varies directly as y and as $(k-y)$, the constant k being the upper limit of growth, or, in other words, the value of the growing variable y at infinite time. Now since the rate of growth of y is dependent upon factors that vary with time, we may generalize (iii) by using $f(x)$ in place of $-a'$, $f(x)$ being some as yet undefined function of time.

Then

$$\frac{dy}{y(k-y)} = f(x)dx.$$

whence

$$\frac{k-y}{my} = e^{-k \int f(x) dx},$$

and

$$y = \frac{k}{1 + me^{-k \int f(x) dx}} = \frac{k}{1 + me^{F(x)}} \quad (\text{iv})$$

where

$$F(x) = -k \int f(x) dx.$$

Then the assumption that the rate of growth of the dependent variable varies as (a) that variable, (b) a constant minus that variable, and (c) an arbitrary function of time, leads to equation (iv), which is of the same form as (i), except that ax has been replaced by $F(x)$. If now we assume that $f(x)$ may be represented by a Taylor series, we have

$$y = \frac{k}{1 + me^{a_1x + a_2x^2 + a_3x^3 + \dots + a_nx^n}} \quad (\text{v})$$

If

$$a_2 = a_3 = a_4 = \dots = a_n = 0$$

then (v) becomes the same as (i).

If m becomes negative the curve becomes discontinuous at finite time. Since this cannot occur in the case of the growth of the organism or of populations, nor indeed so far as we are able to see, for any *phenomenal* changes with time, we shall restrict our further consideration of the equation to positive value only of m . Also since negative values of k would

give negative values of y , which in the case of population or individual growth are unthinkable, we shall limit k to positive values.

With these limitations as to the values of m and k we have the following general facts as to the form of (v). y can never be negative, i.e., less than zero, nor greater than k . Thus the complete curve always falls between the x axis and a line parallel to it at a distance k above it. Further we have the following relations:

$$\begin{aligned} \text{If} \quad F(x) &\doteq \infty & y &= 0 \\ F(x) &\doteq -\infty & y &= k \\ F(x) &\doteq -0 & y &= \frac{k}{1+m} \text{ from below} \\ F(x) &\doteq +0 & y &= \frac{k}{1+m} \text{ from below.} \end{aligned}$$

The maximum and minimum points of (v) occur where $\frac{dy}{dx} = 0$.

$$\text{But } \frac{dy}{dx} = y(k-y) \cdot F'(x),$$

therefore we have maximum and minimum points wherever $F'(x) = 0$.

The fact that $\frac{dy}{dx} = 0$ when either $y = 0$ or $y - k = 0$ shows that the curve passes off to infinity asymptotic to the lines $y = 0$ and $y = k$.

The points of inflection of (v) are determined by the intersections of (v) with the curve.

$$y = \frac{k}{2} - \frac{k}{2} \frac{F''(x)}{[F'(x)]^2} \quad (\text{vi})$$

Since we are seldom justified in using over five arbitrary constants in any practical problem, we may limit equation (v) still further by stopping at the third power of x . This gives the equation

$$y = \frac{k}{1 + me^{a_1x + a_2x^2 + a_3x^3}} \quad (\text{vii})$$

If a_n is positive the curve of equation (v) is reversed and becomes asymptotic to a line AB , at $x = -\infty$ and to the x axis at $x = +\infty$. Thus in equation (vii) a_3 negative is a case of growth, and a_3 positive is a case of decay.

Equation (vii) has several special forms that are of interest, among them being a form similar in shape to the autocatalytic curve (i.e., with no maximum or minimum points and only one point of inflection) except that it is free from the two restrictive features mentioned in our first

paper, that is, location of the point of inflection in the middle and symmetry of the two limbs of the curve. Asymmetrical or skew curves of this sort can only arise when the equation, $F'(x) = 0$, has no real roots. While any odd value of n may yield this form of curve the simplest equation that will do it is that in which $n = 3$, so that the equation of this curve becomes that of (vii).

Having determined that the growth within any one epoch or cycle may be approximately represented by equation (i), or more accurately by (vii), the next question is that of treating several epochs or cycles. Theoretically, some form of (v) may be found by sufficient labor in the adjustment of constants so that one equation with say 5 or 7 constants would describe a long history of growth involving several cycles. Practically, however, we have found it easier and just as satisfactory in other respects to treat each cycle by itself. Since the cycles of any case of growth are additive, we may use for any single cycle the equation

$$y = d + \frac{k}{1 + me^{ka'x}} \quad (\text{viii})$$

or more generally

$$y = d + \frac{k}{1 + me^{a_1x + a_2x^2 + a_3x^3}} \quad (\text{ix})$$

where in both of these forms d represents the total growth attained in all the previous cycles. The term d is therefore the lower asymptote of the cycle of growth under consideration and $d + k$ is its upper asymptote.

In treating any two adjacent cycles, it should be noted that the lower asymptote of the second cycle is frequently below the upper asymptote of the first cycle, due to the fact that the second cycle is often started before the first one has had time to reach its natural level. This for instance would be the case where a population entered upon an industrial era before the country had reached the limit of population possible under purely agricultural conditions.

The theory presented in this paper has been found to be entirely successful in fitting the population growth of many different countries, and in a subsequent publication this fact will be demonstrated with examples.

¹ Papers from the Department of Biometry and Vital Statistics, School of Hygiene and Public Health, The Johns Hopkins University, No. 81.

² Pearl, R. and Reed, L. J., "On the Rate of Growth of the Population of the United States Since 1790 and Its Mathematical Representation." *PROC. NAT. ACAD. SCR.*, Vol. 6, pp. 275-288, 1920.

INDEX

- ABBOT, C. G., FOWLE, F. E., AND ALDRICH L. B. The Larger Results of 20 Years of Solar Radiation Observations, 177.
 Absorption in electric furnace (King), 123.
 Acoustic topography (Barus), 96, 119.
 ADAMS, W. S., AND JOY, A. H. A Method of Deriving the Distance of the A-Type Stars, 173.
 Adrenalin effects upon pigment (Gilson), 130.
 Age distribution, stability of (Lotka), 339.
 Age index of a population (Pearl and LeBlanc), 300.
 ALDRICH, L. B., see Abbot, C. G., 177.
 Aliphatic amines and hydroquinol (Harger), 57.
 Aluminium effects on short X-rays (Duane and Mazumder), 45.
 Amiurus Nebulosus, degeneration and regeneration (Chambers), 186.
 ANDERSON, J. A. The Spectral Energy Distribution and Opacity of Wire Explosion Vapors, 231.
 Animal evolution (Clark), 219.
 Animals, color vision (Hamilton), 350.
 Antiquity of man (Osborn and Reeds), 246.
 Argentite, light effects on (Sheldon and Geiger), 161.
 Atomic sizes (Davis), 61.
 Audition and nerve-deafness (Minton), 274.
 Automobile steering (Webster), 100.
 BACHE FUND, N. A. S., 16.
 Bactericidal properties of radium (Marshall), 317.
 BARUS, CARL. Experiments with the Vacuum Gravitation Needle, 13.
 —. Periods and Logarithmic Decrement of the Gravitation Needle Under High Exhaustion, 63.
 —. Plane Reflection of Sound, as Exhibited by the Pinhole Resonator, 66.
 —. Acoustic Topography Varying with the Position of the Organ Pipe, 96.
 —. Cylindrical Distribution of Nodal Strength Around the Pipe Normal, 119.
 —. Pinhole Resonators, 163.
 —. Static Deflection, Logarithmic Decrement and First Semi-Period of the Vacuum Gravitation Needle, 313.
 —. On a Comparison of the Relative Sensitiveness of Telephones, 326.
 —. The Equilibrium Positions of the Vacuum Gravitation Needle in 1921 and 1922, 327.
 Beryllium and beryllium oxide, structure (McKeehan), 270.
 BLAKESLEE, A. F., AND SINNOTT, E. W. Structural Changes Associated with Factor Mutations and with Chromosome Mutations in *Datura*, 17.
 BLUMBERG, H. New Properties of all Real Functions, 283.
 BRIDGMAN, P. W. The Compressibility of Metals at High Pressures, 361.
 BROWN, E. B. see Johnson, T. B., 187.
 BUNTING, M. A Preliminary Note on Tetramitus, a Stage in the Life Cycle of a Coprozoic Amoeba, 294.
 BURGER, M. H, see Pearl, R., 71.
 CASE, F. H., see Johnson, T. B., 44.
 Chalcedony and silica (Washburn & Navias), 1.
 CHAMBERS, M. H. Degeneration and Regeneration of the Lateral-Line Organs in *Amiurus Nebulosus*, 186.
 Chromosome mutations in *datura* (Sinnot and Blakeslee), 17.
 CLARK, A. H., Animal Evolution, 219.
 CLARK, G. L., AND DUANE, W. A New Method of Using X-Rays in Crystal Analysis, 90.
 CLARKE, F. W., AND WASHINGTON, H. S. The Average Chemical Composition of Igneous Rocks, 108.
 Climatic changes and peat deposits (Dachnowski), 225.
 COBLENTZ, W. W. Effective Temperature of 16 Stars as Estimated from the Energy Distribution in the Complete Spectrum, 49.
 —. Further Measurements of Stellar Temperatures and Planetary Radiation, 330.
 COLE, W. H. Skin Transplantation in Frog Tadpoles, 29.
 Color vision in animals (Hamilton), 350.
 Compressibility of metals (Bridgman), 361.
 Conduction in metals (Hall), 307.
 Coördinates, normal (Veblen), 192.
 Copper, effects on short X-rays (Duane and Mazumder), 45.
 Coprozoic Amoeba (Bunting), 294.
 Coral reefs (Davis), 7.
 Crystal analysis by X-rays (Clark and Duane), 90.
 Crystal structure of beryllium (McKeehan), 270.
 Crystal structure of potassium (McKeehan), 254.
 Cucurbita Pepo seeds, effects on kidney excretion (Masurovsky), 39.

- Curve, continuous (Moore), 33.
 Cyanosis, certain factors involved in (Lundsgaard and Van Slyke), 280.
- DACHNOWSKI, A. P. The Correlation of the Units and Climatic Changes in Peat Deposits of the United States and Europe, 225.
 Dark nebulae (Russell), 115.
 Datura, mutations and structure (Sinnott and Blakeslee), 17.
 DAVIS, B. Ionization and Radiation Potentials and Size of the Atom, 61.
 DAVIS, B., AND TERRILL, H. M. Grating Space of Mica and the Intensities of Spectral Orders, 59.
 —. The Refraction of X-rays in Calcite, 357.
 DAVIS, W. M. Coral Reefs of the Louisiade Archipelago, 7.
 Deaf, aids for the (Wegel), 155.
 Deafness and theories of audition (Minton), 274.
 Density of population (Pearl and Parker), 212.
 Diazomethane, action on uracil (Johnson, Hill and Case), 44.
 Distance of A-type stars (Adams and Joy), 173.
 Drosophila, reproduction in (Pearl and Parker), 212.
 DUANE, W., see Clark, G. L., 90.
 DUANE, W., AND MAZUMDER, K. C. Absorption of Short X-rays by Aluminium and Copper, 45.
 DUANE, W., AND PATTERSON, R. A. Note on X-ray Spectra, 85.
- Ear, sensitivity of (Fletcher and Wegel), 5; (Mackenzie), 188.
 Einstein theory (Eisenhart), 24.
 EISENHART, L. P. Ricci's Principal Directions for a Riemann Space and the Einstein Theory, 24.
 —. Fields of Parallel Vectors in the Geometry of Paths, 207.
 —. Spaces with Corresponding Paths, 233.
 EISENHART, L. P., AND VELEN, O. The Riemann Geometry and its Generalization, 19.
 ELDRIDGE, J. A. The Energy Losses Accompanying Ionization and Resonance in Mercury Vapor, 268.
 Electric furnace absorption (King), 123.
 Electromagnetic theory (Lorentz), 333.
 Electron theory conduction (Hall), 307.
 Electronic bombardment of nickel (Farnsworth), 251.
 Energetics of evolution (Lotka), 147.
 Energy distribution in stellar spectra (Coblentz), 49.
 EPSTEIN, P. S. The Evaluation of Quantum Integrals, 166.
- Ether, effect on pigment (Wyman), 128.
 EVANS, H. M., AND LONG, J. A. Characteristic Effects upon Growth, Oestrus and Ovulation, Induced by the Intra-peritoneal Administration of Fresh Anterior Hypophyseal Substance, 38.
 Evolution, animal (Clark), 219.
 Evolution, arrested (Ruedemann), 54.
 Evolution, energetics of (Lotka), 147.
 Explosion vapors, spectra of (Anderson), 231.
- FARNSWORTH, H. E. Electronic Bombardment of Nickel, 251.
 FISCHER, C. A. Note on the Definition of a Lineal Functional, 26.
 FLETCHER, H., AND WEGEL, R. L. Frequency-Sensitivity of Normal Ears, 5.
 Fossils of New York silurian (Ruedemann), 55.
 FOWLE, F. E., see Abbot, C. C., 177.
 FRANKLIN, P. The Meaning of Rotation in the Special Theory of Relativity, 265.
 Frequency-sensitivity of ears (Fletcher and Wegel), 5.
 Functional, lineal (Fischer), 26.
 Functions, properties of real (Blumberg), 283.
- Gaseous elements at high temperatures (Noyes and Wilson), 303.
 GEIGER, P. H., see Sheldon, H. H., 161.
 General analysis (Moore), 288.
 Geometry of paths (Veblen), 192, 347; (Eisenhart), 207, 233.
 Geometry, Riemann, generalized (Eisenhart and Veblen), 19.
 GILSON, A. S. JR. The Diverse Effects of Adrenalin upon the Migration of the Scale and Retinal Pigment in the Fish, *Fundulus Heteroclitus*, Linn, 130.
 Grating space of mica (Davis and Terrill), 59.
 Gravitational field of sun (Synge), 204.
 Gravitational needle (Barus), 13, 63, 313, 327.
 Group of substitutions (Miller), 238.
 Growth effects of hypophyseal substance (Evans and Long), 38.
- HALE, G. E. Invisible Sun-Spots, 168.
 HALL, E. H. An Electron Theory of Electric Conduction in Metals, 307.
 HALSTED, W. S. Replantation of Entire Limbs without Suture of Vessels, 181.
 HAMILTON, W. F. A Direct Method of Testing Color Vision in Lower Animals, 650.
 HARGER, R. N. The Oxidation of Hydroquinol in Presence of Aliphatic Amines, 57.

- HARRISON, G. R. The Absorption of Light by Sodium and Potassium Vapors, 260.
- Hearing, aids for the deaf (Wegel), 155.
- Hearing, sensitivity of (Fletcher and Wegel), 5; (MacKenzie), 188.
- Heaviside's theorem (Lorentz), 333.
- Hemoglobin concentration and cyanosis (Lundsgaard and Van Slyke), 280.
- Hesperopithecus (Osborn), 245.
- HILL, A. J., see Johnson, T. B., 44.
- HITCHCOCK, F. L. A Solution of Linear Matrix Equation by Double Multiplication, 78.
- HOWARD, L. O. A Side Line in the Importation of Insect Parasites of Injurious Insects from One Country to Another, 133.
- Hydroquinol, oxidation of, with aliphatic amines (Harger), 57.
- Hypophyseal substance affecting growth (Evans and Long), 38.
- Igneous rocks, analysis of (Clarke and Washington), 108.
- Insects parasitic on others (Howard), 133.
- Insects, photic orientation (Mast), 240.
- Integrals, quantum (Epstein), 166.
- Invisible sun-spots (Hale), 168.
- Ionization at high temperatures (Noyes and Wilson), 303.
- Ionization in electric furnace (King), 123.
- Ionization in mercury vapor (Eldridge), 268.
- Ionization of potassium vapor (Williamson), 255.
- Ionization potentials (Davis), 61.
- Jades of Middle America (Washington), 319.
- JOHNSON, T. B., AND BROWN, E. B. The Isolation of Pyrimidines from the Nucleic Acid of Tubercle Bacillus, 187.
- JOHNSON, T. B., HILL, A. J., AND CASE, F. H. The Action of Diazomethane on Uracil, 44.
- JOY, A. H., see Adams, W. S., 173.
- Jupiter, perturbations of minor planets (Leuschner), 170.
- Kidney excretion, as effected by Cucurbita Pepo seeds (Masurovsky), 39.
- KING, A. S. Ionization and Absorption Effects in the Electric Furnace, 123.
- LÉBLANC, T. J., see Pearl, R., 300.
- LEUSCHNER, A. O. Comparison of Theory with Observation for the Minor Planets 10 Hygiea and 175 Andromache with Respect to Perturbations by Jupiter, 170.
- Light effects on argentite (Sheldon and Geiger), 161.
- Limbs, replantation of (Halsted), 181.
- Limits, generalized (Moore), 288.
- Lineal functional (Fischer), 26.
- Linear matrix equation (Hitchcock), 78.
- LONG, J. A., see Evans, H. M., 38.
- LORENTZ, H. A. Proof of Theorem due to Heaviside, 333.
- LOTKA, A. J. Contribution to the Energetics of Evolution, 147.
- Natural Selection as a Physical Principle, 151.
- The Stability of the Normal Age Distribution, 339.
- Louisiade Archipelago (Davis), 7.
- LUNDGAARD, C., AND VAN SLYKE, D. D. The Quantitative Influences of Certain Factors Involved in the Production of Cyanosis, 280.
- McKREHAN, L. W. The Crystal Structure of Beryllium and of Beryllium Oxide, 270.
- The Crystal Structure of Potassium, 254.
- MACKENZIE, D. The Relative-Sensitivity of the Ear at Different Levels of Loudness, 188.
- Magellanic cloud, distance of (Shapley), 69.
- Man, antiquity of (Osborn and Reeds), 246.
- MARSHALL, J. A. Bactericidal Properties of the Products of Radium Emanation, 317.
- MAST, S. O. Photic Orientation in Insects, 240.
- MASUROVSKY, B. A Study of the Effects of Cucurbita Seeds on Kidney Excretion, 39.
- Matrix equations (Hitchcock), 78.
- MAZUMDER, K. C., see Duane, W., 45.
- Measurements, theory of (Shewhart), 248.
- Mercury vapor, ionization of (Eldridge), 268.
- Metallic conduction (Hall), 307.
- Metals, compressibility (Bridgman), 361.
- Mica, grating space (Davis and Terrill), 59.
- Middle American Jades (Washington), 319.
- MILLER, G. A. Number of Substitutions Omitting at Least One Letter in a Transitive Group, 238.
- MINER, J. R. The Probable Error of the Vital Index of a Population, 106.
- MINTON, J. P. Some Cases of Nerve-Deafness and their Bearing on Resonance Theories of Audition, 274.
- MOORE, C. N. Generalized Limits in General Analysis, 288.

- MOORE, R. L. On the Relation of a Continuous Curve to its Complementary Domains in Space of Three Dimensions, 33.
- Mutations in *datura* (Sinnott and Blakeslee), 17.
- NATIONAL ACADEMY OF SCIENCES, Bache Fund, 16.
- Natural selection as a physical principle (Lotka), 151.
- NAVIAS, L., AND E. W. WASHBURN. The Relation of Chalcedony to the Other Forms of Silica, 1.
- Nebulae, dark (Russell), 115.
- Needle, gravitation (Barus), 13, 63, 313, 327.
- Nerve deafness (Minton), 274.
- NETTLETON, L. L. Characteristics of a Short Wave Oscillator at Very Low Pressures, 353.
- Nickel, electronic bombardment of (Farnsworth), 251.
- NOYES, A., AND WILSON, H. A. The Thermal Ionization of Gaseous Elements at High Temperatures, 303.
- Orientation, photic (Mast), 240.
- OSBORN, H. F. *Hesperopithecus*, the First Anthropid Primate Found in America, 245.
- OSBORN, H. F., AND REEDS, C. A. Recent Discoveries on the Antiquity of Man, 246.
- Oscillator, short wave (Nettleton), 353.
- Parallel vectors in geometry (Eisenhart), 207.
- Parasites of injurious insects (Howard), 133.
- PARKER, S. L., see Pearl, R., 212.
- Paths, geometry of (Veblen), 192, 347; (Eisenhart), 207, 233.
- PATTERSON, R. A., see Duane, W., 85.
- PEARL, R. Seasonal Fluctuations of the Vital Index of a Population, 76.
- PEARL, R. AND BACON, A. L. Biometrical Study of the Relations of Certain Viscera in Tuberculosis, 125.
- PEARL, R., AND BURGER, M. H. The Vital Index of Population of England and Wales, 1838-1920, 71.
- PEARL, R., AND LeBLANC, T. J. A Further Note on the Age Index of a Population, 300.
- PEARL, R., AND PARKER, S. L. On the Influence of Density of Population upon the Rate of Reproduction in *Drosophila*, 212.
- PEARL, R. AND REED, L. J. A Further Note on the Mathematical Theory of Population Growth, 365.
- Peat deposits and climatic changes (Dachnowski), 225.
- Persistence of animals (Ruedemann), 54.
- Perturbations of minor planets (Leuschner), 170.
- Physical quantity, measurement of (Shewhart), 248.
- Pigment in fish (Wyman), 128; (Gilson), 130.
- Pinhole resonators (Barus), 66, 96, 119, 163.
- Planets, minor (Leuschner), 170.
- Polynomial, roots of the derivative of, (Walsh), 139.
- Population, age distribution (Lotka), 339.
- Population, age index of (Pearl and LeBlanc), 300.
- Population, growth of (Pearl and Reed), 365.
- Population, vital index of (Pearl and Berger), 71; (Pearl), 76; (Miner), 106.
- Potassium, crystal structure (McKeehan), 254.
- Potassium vapor, the ionization of (Williamson), 255.
- Potassium vapors, absorption by (Harrison), 260.
- Principal directions in space (Eisenhart), 24; (Synge), 198, 204.
- Pyrimidines from tubercle bacillus (Johnson and Brown), 187.
- Quantum integrals (Epstein), 166.
- Radiation of the sun (Abbot, Fowle and Aldrich), 177.
- Radiation potentials (Davis), 61.
- Radium emanation, bactericidal properties (Marshall), 317.
- Real functions (Blumberg), 283.
- REED, L. J. See Pearl, R., 365.
- REEDS, C. A., see Osborn, H. F., 246.
- Refraction of X-rays (Davis and Terrill), 357.
- Regeneration in *Amiurus* (Chambers), 186.
- Relativity, rotation in (Franklin), 265.
- Replantation of limbs (Halsted), 181.
- Reproduction and density of population (Pearl and Parker), 212.
- Resonators, pinhole (Barus), 66, 96, 119, 163.
- Ricci's principal directions (Eisenhart), 24.
- Riemann geometry generalized (Eisenhart and Veblen), 19.
- Rocks, chemical composition of (Clarke and Washington), 108.
- Roots of polynomial derivatives (Walsh), 139.
- Rotation in relativity (Franklin), 265.
- RUEDEMANN, R. Additional Studies in Arrested Evolution, 54.
- . New Forms of Life from the Silurian, 55.
- RUSSELL, H. N. Dark Nebulae, 115.

- Sensitivity of the ear (Fletcher and Wegel), 5; (MacKenzie), 188.
 Sensitivity of Telephones (Barus), 326.
 SHAPLEY, H. Note on Problem of Great Stellar Distances, 69.
 SHELDON, H. H., AND GEIGER, P. H. The Production of an E. M. F. on Closed Circuit by a Light Effect on Argentite, 161.
 SHEWHART, W. A. On the Measurement of a Physical Quantity Whose Magnitude is Influenced by Primary Causes beyond the Control of the Observer and on the Method of Determining the Relation between Two Such Quantities, 248.
 Silica and chalcedony (Washburn and Navias), 1.
 Silurian life (Ruedemann), 54.
 SINNOTT, E. W., see Blakeslee, A. F., 17.
 Skin, transplantation (Cole), 29.
 Sodium vapors, absorption by (Harrison), 260.
 Solar radiation (Abbot, Fowle and Aldrich), 177.
 Sound, reflection of (Barus), 66.
 Space, principal direction (Eisenhart), 24; (Synge), 198, 204.
 Space curves (Moore), 33.
 Spaces with corresponding paths (Eisenhart), 233.
 Spectra of explosion vapors (Anderson), 231.
 Spectra of X-rays (Duane and Patterson), 85.
 Stars, distance of (Adams and Joy), 173.
 Stars, distribution of velocities (Strömberg), 141.
 Stars, temperatures (Coblentz), 49, 330.
 Steering around a corner (Webster), 100.
 Stellar distances (Shapley), 69.
 STRÖMBERG, G. On the Distribution of the Velocities of Stars of Late Types of Spectrum, 141.
 Substitution groups (Miller), 238.
 Sun-spots, invisible (Hale), 168.
 SYNGE, J. L. Principal Directions in a Riemannian Space, 198.
 —. Principal Directions in the Einstein Solar Field, 204.
 Tadpoles, skin transplantation (Cole), 29.
 Telephones, relative sensitiveness (Barus), 326.
 Temperature of stars (Coblentz), 49, 330.
 TERRILL, H. M., see Davis, B., 59.
 —. See Davis, B., 357.
 Tetramitus (Bunting), 294.
 Time units in paleontology (Dachnowski), 225.
 Transplantation of skin in tadpoles (Cole), 29.
 Tubercle bacillus, pyrimidines from (Johnson and Brown), 187.
 Tuberculosis and viscera (Pearl and Bacon), 125.
 Uracil, as effected by diazomethane (Johnson, Hill and Case), 44.
 Vacuum gravitation needle (Barus), 13, 63, 313, 327.
 Vacuum tubes (Nettleton), 353.
 VAN SLYKE, D. D., see Lundsgaard, C., 280.
 VEBLEN, O. Normal Coördinates for the Geometry of Paths, 192.
 —. Projective and Affine Geometry of Paths, 347.
 —. See Eisenhart, L. P., 19.
 Velocity distribution for stars (Strömberg), 141.
 Viscera in tuberculosis (Pearl and Bacon), 125.
 Vital index (Pearl and Burger), 71; (Pearl), 76; (Miner), 106.
 WALSH, J. L. On the Location of the Roots of the Derivative of a Polynomial, 139.
 WASHBURN, E. W., see Navias, L., 1.
 WASHINGTON, H. S. The Jades of Middle America, 319.
 —. See Clarke, F. W., 108.
 WEBSTER, A. G. On Steering an Automobile around a Corner, 100.
 WEGEL, R. L. The Physical Examination of Hearing and Binaural Aids for the Deaf, 155.
 —. See Fletcher, H., 5.
 WILLIAMSON, R. C. Preliminary Report on the Ionization Vapor of Potassium by Light, 255.
 WILSON, H. A., see Noyes, A., 303.
 WYMAN, L. C. The Effect of Ether upon the Migration of the Scale Pigment and the Retinal Pigment in the Fish, *Fundulus Heteroclitus*, 128.
 X-ray crystal analysis (Clark and Duane), 90.
 X-ray spectra (Duane and Patterson), 85.
 X-rays, absorbed by aluminium and copper (Duane and Mazumder), 45.
 X-rays in calcite (Davis and Terrill), 357.

INFORMATION TO CONTRIBUTORS

THE PROCEEDINGS is the official organ of the NATIONAL ACADEMY OF SCIENCES and of the NATIONAL RESEARCH COUNCIL for the publication of brief accounts of important current researches of members of the ACADEMY and of the COUNCIL and of other American investigators. The Proceedings will aim especially to secure prompt publication of original announcements of discoveries and wide circulation of the results of American research among investigators in other countries and in all branches of science.

ARTICLES should be brief. The viewpoint should be comprehensive in giving the relation of the paper to previous publications of the author or of others and in exhibiting, where practicable, the significance of the work for other branches of science. Elaborate technical details of the work and long tables of data should be avoided, but authors should be precise in making clear the new results and should give some record of the methods and data upon which they are based.

MANUSCRIPTS should be prepared with a current number of the Proceedings as a model in matters of form, and should be typewritten in duplicate with double spacing, the author retaining one copy. Illustrations should be confined to text-figures of simple character, though more elaborate illustrations may be allowed in special instances to authors willing to pay for their preparation and insertion. Particular attention should be given to arranging tabular matter in a simple and concise manner.

REFERENCES to literature, numbered consecutively, will be placed at the end of the article and short footnotes should be avoided. It is suggested that references to periodicals be furnished in some detail and in general in accordance with the standard adopted for the Subject Catalogue of the International Catalogue of Scientific Literature, viz., name of author, with initials following (ordinarily omitting title of paper), abbreviated name of Journal, with place of publication, series (if any), volume, year, inclusive pages. For example: Montgomery, T. H., *J. Morph., Boston*, 22, 1911 (731-815); or, Wheeler, W. M., *Königsburg, Schr. physik. Ges.*, 55, 1914 (1-142).

PAPERS by members of the Academy may be sent to Edwin Bidwell Wilson, Managing Editor, Harvard School of Public Health, Boston 17, Mass. Papers by non-members of the ACADEMY or COUNCIL should be submitted through some member.

PROOF will not ordinarily be sent; if an author ask for proof, it will be sent with the understanding that charges for his corrections shall be billed to him. Authors are therefore requested to make final revisions on the typewritten manuscripts. The editors cannot undertake to do more than correct obvious minor errors.

REPRINTS should be ordered at the time of submission of manuscript. They will be furnished to authors at cost, approximately as follows:

SCHEDULE OF RATES FOR REPRINTS, CARRIAGE CHARGES EXTRA.					
No. copies	4 pp.	8 pp.	12 pp.	16 pp.	Covers
50	\$1.95	\$3.90	\$5.85	\$7.80	\$1.50
100	2.22	4.44	6.66	8.88	1.89
150	2.49	4.98	7.47	9.96	2.27
200	2.76	5.52	8.28	11.04	2.66
250	3.03	6.06	9.09	12.12	3.05

Copyright, 1922, by the National Academy of Sciences

CONTENTS

	Page
MATHEMATICS.—PROJECTIVE AND AFFINE GEOMETRY OF PATHS <i>By O. Veblen</i>	347
PHYSIOLOGY.—A DIRECT METHOD OF TESTING COLOR VISION IN LOWER ANIMALS <i>By W. F. Hamilton</i>	350
PHYSICS.—CHARACTERISTICS OF A SHORT WAVE OSCILLATOR AT VERY LOW PRESSURES <i>By L. L. Nettleton</i>	353
PHYSICS.—THE REFRACTION OF X-RAYS IN CALCITE	
<i>By Bergen Davis and H. M. Terrill</i>	357
PHYSICS.—THE COMPRESSIBILITY OF METALS AT HIGH PRESSURES	
<i>By P. W. Bridgman</i>	361
STATISTICS.—A FURTHER NOTE ON THE MATHEMATICAL THEORY OF POPULATION GROWTH <i>By Raymond Pearl and Lowell J. Reed</i>	365
INDEX	369



